

## *A Study in Step Size*

*Temple H. Fay*

*University of Southern Mississippi*

*Hattiesburg, MS 39406-5045*

*Mathematics Magazine*, April 1997, Volume 70, Number 2, pp.116–117.

*Mathematics Magazine* is an official publication of the Mathematical Association of America

While experimenting with giving polar plots some “texture,” I discovered an interesting effect that might intrigue students: high sensitivity to step size. The base equation is that used to generate the butterfly curve (see [1], for example)

$$r(\theta) = e^{\cos(\theta)} - 2 \cos(4\theta),$$

but the technique can be applied, and the same effect observed, using almost any polar equation.

The “texture” is obtained by multiplying this base curve by a rapidly varying sinusoidal factor, in this case by  $\sin^4(\lambda\theta)$ , where  $\lambda = 99999999$ . The fourth power was chosen to keep the factor non-negative and small. The value of  $\lambda$  was chosen arbitrarily; any large number would produce the same effect.

Data sets, consisting of points  $(x_n, y_n)$  where

$$\rho(\theta) = (e^{\cos(\theta)} - 2 \cos(4\theta))\sin^4(\lambda\theta)$$

and

$$x_n = \rho(\theta_n)\sin(\theta_n)$$

$$y_n = \rho(\theta_n)\cos(\theta_n),$$

were produced by setting  $\theta_0 = 0$ , and  $\theta_n = \theta_{n-1} + h$  where  $h$  denotes the step size, for  $0 \leq \theta < 2\pi$ . (Reversing the sine and cosine from their “usual” positions rotates the butterfly  $90^\circ$  into the upright position shown.) These data sets contain roughly 11,500 to 42,000 points.

The plots shown on the cover of this issue were produced with step sizes as follows:


$$h_1 = 0.00015 \quad h_2 = 0.0003$$

$$h_3 = 0.0005 \quad h_4 = 0.00055$$

The plots in the following Figure used these step sizes:

$$h_5 = 0.0007 \quad h_6 = 0.000169$$

$$h_7 = 0.000711 \quad h_8 = 0.00071$$



**Figure**  
Four butterfly curves

Watching the dynamics of the plotting of the sequentially-generated points  $(x_n, y_n)$  is interesting in and of itself. Students might enjoy experimenting with different equations, values of  $\lambda$ , and step sizes.

REFERENCE

1. T. H. Fay, The butterfly curve, *American Mathematical Monthly* 96 (1989), 442–443.