


$n = 2$ , the solution is  $x = y = u - 1 = z - 2$ . The generalization of equations (1) and (2) can continue, but as Fermat so aptly put it, the margin is too small.



Other cases of Fermat's craniosis have recently come to public attention. It is alarming that a preponderance of these documented cases have occurred within the mathematical community. In some cases, the infection has been traced back 400 years to an association with Fermat himself. The disease is eventually fatal, and the most debilitating effect is the patient's zealous desire to study point-set topology.

The good news is that only number theorists are susceptible to infection.



## Behold! The Midpoint Rule is Better Than the Trapezoidal Rule for Concave Functions

Frank Buck, California State University, Chico, CA



## The Bisection Algorithm is Not Linearly Convergent

Sui-Sun Cheng and Tzon-Tzer Lu, Tsing Hua University, Taiwan, Republic of China

The bisection algorithm is a method for finding a root of the equation  $f(x) = 0$ , where  $f$  is continuous on  $[a, b]$  and  $f(a)f(b) < 0$ . Let  $a(1) = a$ ,  $b(1) = b$  and let  $p(1)$  be the midpoint of  $[a(1), b(1)]$ . Now let  $[a(2), b(2)]$  be one of the halves of this interval for which  $f[a(1)]f[b(2)] < 0$ , and let  $p(2)$  be the midpoint of  $[a(2), b(2)]$ . Proceeding in this manner, the bisection algorithm generates a sequence of bracketing intervals  $[a(i), b(i)]$  and a sequence of midpoints  $p(i)$  of these intervals. It is well known that  $p(i)$  converges to a root  $p$  of  $f$ . Some authors go even further and assert that  $p(i)$  converges to  $p$  linearly—that is, there is some integer  $N$  and some constant  $Q$  such that  $|p(i+1) - p| \leq Q|p(i) - p|$  for all  $i \geq N$ . This is not true in general. Although other authors have not made such an assertion, it is surprising that they give no counterexample. Here is a counterexample.

Let  $p = 2^{-1} + 2^{-4} + \dots + 2^{-n^2} + \dots$  and let  $f(x) = x - p$ . Let the first bracketing interval of the root  $p$  be  $[2^{-1}, 1]$ . Then it is easily verified that the second and the third bracketing intervals are  $[2^{-1}, 2^{-1} + 2^{-2}]$  and  $[2^{-1}, 2^{-1} + 2^{-3}]$ . In general, let  $S(n) = \{n^2, n^2 + 1, \dots, (n+1)^2 - 1\}$  for  $n = 1, 2, \dots$ . Then these sets form a partition of the set of positive integers. We assert that for  $i \in S(n)$ , the  $i$ th bracketing interval is

$$[a(i), b(i)] = [2^{-1} + 2^{-4} + \dots + 2^{-n^2}, 2^{-1} + 2^{-4} + \dots + 2^{-n^2} + 2^{-i}]. \quad (*)$$

The proof is by induction. We have seen above that  $(*)$  holds for each  $i \in S(1)$ .