

# Partial Fraction Decomposition by Division

Sidney H. Kung (sidneykung@yahoo.com) University of North Florida, Jacksonville FL 32224

In this note, we present a method for the partial fraction decomposition of two algebraic functions: (i)  $f(x)/(ax + b)^t$  and (ii)  $f(x)/(px^2 + qx + r)^t$ , where  $f(x)$  is a polynomial of degree  $n$ ,  $t$  is a positive integer, and  $px^2 + qx + r$  is an irreducible ( $q^2 < 4pr$ ) quadratic polynomial. Our algorithm is relatively simple in comparison with those given elsewhere [1, 2, 3, 4, 5, 6, 7, 8]. The essence of the method is to use repeated division to re-express the numerator polynomial in powers of the normalized denominator. Then upon further divisions, we obtain a sum of partial fractions in the form  $A_i/(ax + b)^i$  or  $(B_jx + C_j)/(px^2 + qx + r)^j$  for the original function.

For (i), we let  $c = b/a$ , and express  $f(x)$  as follows:

$$f(x) = A_n(x + c)^n + A_{n-1}(x + c)^{n-1} + \dots + A_2(x + c)^2 + A_1(x + c) + A_0$$

$$= (x + c)[A_n(x + c)^{n-1} + A_{n-1}(x + c)^{n-2} + \dots + A_2(x + c) + A_1] + A_0,$$

where each  $A_i$  is a real coefficient to be determined. Then the remainder after we divide  $f(x)$  by  $x + c$  gives the value of  $A_0$ . The quotient is

$$q_1(x) = (x + c)[A_n(x + c)^{n-2} + A_{n-1}(x + c)^{n-3} + \dots + A_3(x + c) + A_2] + A_1.$$

If we now divide  $q_1(x)$  by  $x + c$ , we see that the next remainder is  $A_1$  and the quotient is

$$q_2(x) = (x + c)[A_n(x + c)^{n-3} + A_{n-1}(x + c)^{n-4} + \dots + A_3] + A_2.$$

Continuing to divide in this manner  $n - 1$  times, we get the quotient  $q_{n-1}(x) = A_n(x + c) + A_{n-1}$ . Finally, dividing  $q_{n-1}(x)$  by  $x + c$ , we obtain the last two coefficients,  $A_{n-1}$  and  $A_n$ . Thus, it follows that

$$\frac{f(x)}{(ax + b)^t} = \frac{1}{a^t} \left[ \frac{A_n}{(x + c)^{t-n}} + \frac{A_{n-1}}{(x + c)^{t-n+1}} + \dots + \frac{A_1}{(x + c)^{t-1}} + \frac{A_0}{(x + c)^t} \right]. \quad (1)$$

For example, to find the partial fraction decomposition of  $(x^4 + 2x^3 - x^2 + 5)/(2x - 1)^5$ , we use  $c = -1/2$  and perform synthetic division to obtain  $A_0$  through  $A_n$ .

1/2)	1	2	-1	0	5	
		1/2	5/4	1/8	1/16	
	1	5/2	1/4	1/8		81/16 ← A <sub>0</sub>
		1/2	3/2	7/8		
	1	3	7/4		1	← A <sub>1</sub>
		1/2	7/4			
	1	7/2		7/2		← A <sub>2</sub>
		1/2				
A <sub>4</sub> ⇒	1	4				← A <sub>3</sub>

Substituting the coefficients into (1), we have

$$\begin{aligned} \frac{x^4 + 2x^3 - x^2 + 5}{(2x - 1)^5} &= \frac{1}{2^5} \left[ \frac{1}{(x - 1/2)} + \frac{4}{(x - 1/2)^2} + \frac{7/2}{(x - 1/2)^3} + \frac{1}{(x - 1/2)^4} \right. \\ &\quad \left. + \frac{81/16}{(x - 1/2)^5} \right] \\ &= \frac{1/16}{2x - 1} + \frac{1/2}{(2x - 1)^2} + \frac{7/8}{(2x - 1)^3} \\ &\quad + \frac{1/2}{(2x - 1)^4} + \frac{81/16}{(2x - 1)^5}. \end{aligned}$$

For (ii), we let  $u = q/p$ ,  $v = r/p$ , and express  $f(x)$  in the following form:

$$\begin{aligned} f(x) &= B_{(n-1)/2}(x^2 + ux + v)^{(n-1)/2} + B_{(n-3)/2}(x^2 + ux + v)^{(n-3)/2} + \dots \\ &\quad + B_1(x^2 + ux + v) + B_0, \end{aligned}$$

where each coefficient  $B_k$ ,  $k = 0, 1, \dots, (n-1)/2$ , is a linear function of  $x$ , and where we assume that  $n \leq 2t - 1$ . In this case, dividing  $f(x)$  and each successive quotient by  $x^2 + ux + v$  as described above, we obtain

$$\begin{aligned} \frac{f(x)}{(px^2 + qx + r)^t} &= \frac{1}{p^t} \left[ \frac{B_{(n-1)/2}}{(x^2 + ux + v)^{t-(n-1)/2}} \right. \\ &\quad \left. + \frac{B_{(n-3)/2}}{(x^2 + ux + v)^{t-(n-3)/2}} + \dots + \frac{B_0}{(x^2 + ux + v)^t} \right]. \end{aligned} \quad (2)$$

For instance, take the rational function  $(x^5 - 4x^4 + 3x^2 - 2)/(x^2 - x + 2)^3$ . Then  $u = -1$  and  $v = 2$ . Since  $(n-1)/2 = 2$ , we let

$$x^5 - 4x^4 + 3x^2 - 2 = (Mx + N)(x^2 - x + 2)^2 + (Kx + L)(x^2 - x + 2) + Ix + J.$$

Since most students are not familiar with the synthetic division technique when the divisor is a quadratic polynomial, long division can be used in place of the following computation to find the coefficients  $I, J, K, L, M$ , and  $N$ .

$$\begin{array}{r} \begin{array}{cccccc|cccc} 1 & -4 & 0 & 3 & 0 & -2 & 1 & -3 & -5 & 4 \\ 1 & -1 & 2 & & & & 1 & -1 & 2 & \end{array} \\ \hline \begin{array}{cccc} -3 & -2 & 3 & \\ -3 & 3 & 6 & \end{array} \\ \hline \begin{array}{cccc} -5 & 9 & 0 & \\ -5 & 5 & -10 & \end{array} \\ \hline \begin{array}{ccc} 4 & 10 & -2 \\ 4 & -4 & 8 \end{array} \\ \hline \begin{array}{cc} 14 & -10 \\ I & J \end{array} \end{array}$$

$$\begin{array}{cccc|cc}
 & & & & M & N \\
 1 & -3 & -5 & 4 & 1 & -2 \\
 1 & -1 & 2 & & 1 & -1 & 2 \\
 \hline
 & -2 & -7 & 4 & & & \\
 & -2 & 2 & -4 & & & \\
 \hline
 & & -9 & 8 & & & \\
 & & K & L & & & 
 \end{array}$$

Substituting the coefficients in (2) (note that  $t - (n - 1)/2 = 1$ ) gives

$$\frac{x^5 - 4x^4 + 3x^2 - 2}{(x^2 - x + 2)^3} = \frac{x - 2}{x^2 - x + 2} + \frac{-9x + 8}{(x^2 - x + 2)^2} + \frac{14x - 10}{(x^2 - x + 2)^3}.$$

Note also that  $x^2 - x + 2 = (x - 1/2)^2 + 7/4$ . On the right hand side of the above expression, replacing the coefficients  $-2$ ,  $8$ , and  $-10$  in the numerators by  $-2 + 1/2M$ ,  $8 + 1/2M$ , and  $-10 + 1/2M$ , respectively, we get

$$\begin{aligned}
 \frac{x^5 - 4x^4 + 3x^2 - 2}{((x - 1/2)^2 + 7/4)^3} &= \frac{(x - 1/2) - 3/2}{(x - 1/2)^2 + 7/4} + \frac{-9(x - 1/2) + 7/4}{((x - 1/2)^2 + 7/4)^2} \\
 &+ \frac{14(x - 1/2)^2 - 3}{((x - 1/2)^2 + 7/4)^3}.
 \end{aligned}$$

This last expression is an easily antiderivable form.

## References

1. S. Burgstahler, An alternative for certain partial fractions, *College Math. J.* **15** (1984) 57–58.
2. X.-C. Huang, A short cut to partial fractions, *College Math. J.* **22** (1991) 413–415.
3. P. T. Joshi, Efficient techniques for partial fractions, *College Math. J.* **14** (1983) 110–118.
4. J. E. Nymann, An alternative for partial fractions (part of the time), *College Math. J.* **14** (1983) 60–61.
5. P. Schultz, An algebraic approach to partial fractions, *College Math. J.* **14** (1983) 346–348.
6. M. R. Spiegel, Partial fractions with repeated linear or quadratic factors, *Amer. Math. Monthly* **57** (1950) 180–181.
7. T. N. Subramaniam and D. E. G. Malm, How to integrate rational functions, *Amer. Math. Monthly* **99** (1992) 762–772.
8. J. Wiener, An algebraic approach to partial fractions, *College Math. J.* **17** (1986) 71–72.

————— o —————

## An Elegant Mode for Determining the Mode

D.S. Broca (dsbroca@iimk.ac.in), Indian Institute of Management Kozhikode, Kozhikode 673 571, Kerala, India

For any probability distribution, the mode, like the mean and median, is a measure of central tendency. Geometrically, it represents the relative maximum of the probability density function (pdf) and thus is the most striking feature in the curve's topogra-