

# The Logarithm of $-1$ \*

Dominic Klyve<sup>†</sup>

August 7, 2023

## 1 Introduction

Logarithms were introduced to the world by John Napier (1550–1617) in 1614, in a work entitled *A Description of the Wonderful Table of Logarithms* (or, in the Latin original, *Mirifici logarithmorum canonis descriptio*), as a bridge between an arithmetic sequence and a geometric one. Four hundred years later, logarithms are ubiquitous in mathematics and science, and it is difficult to imagine a world without them.

The eighteenth century was in a time of transition in the story of logarithms (as it was in so many fields of mathematics). The elementary properties of logs were well understood, but many questions about them remained. This project will focus on one of these questions—namely, how to extend the domain of the logarithm function to negative numbers.

Recall that  $\log_b(x) = a$  is equivalent to  $b^a = x$ , so, for example,  $\log_{10}(1000) = 3$ . Given any function, it is convenient for its domain to be as large as possible, and for mathematical reasons (the applications would come many years later), eighteenth-century mathematicians wished to find a way to do this for the logarithm function. As it turns out, deciding the best way to do so was not easy, and would lead to a serious disagreement in the mathematical community.

**Task 1** Why is it difficult to find the logarithm of a negative number? For example, why can't you easily find a value for  $\log_{10}(-100)$ ?

## 2 Functions with Several Values?

You have likely seen a formal definition of a function in the past. Although there are small differences between texts, the definition usually looks something like this:

**Definition.** Let  $X$  and  $Y$  be sets. A **function**  $f : X \rightarrow Y$  defined on  $X$  is a rule that assigns to each element  $x \in X$  an element  $y \in Y$ , in which case we write  $y = f(x)$ .

**Task 2** Check the definition of function in a current textbook. Does it differ in any significant way from the definition above? If so, how?

---


\*This updated version of the project replaces the version that was dated June 17, 2021 and published in *Convergence* in June 2021. The author is grateful to Robert Bradley, whose article “Euler, D’Alembert and the Logarithm Function” [Bradley, 2007] provided the central ideas for much of this project.

<sup>†</sup>Department of Mathematics, Central Washington University, Ellensburg, WA 98926; dominic.klyve@cwu.edu.

**Task 3** A very important property of functions seems to be that for each  $x$ ,  $f(x)$  has only one value. (This is related to the “vertical line test” that you have seen in calculus.) Can you think of any “function” (possibly expanding or modifying the definition a bit) for which  $f(x)$  has more than one value? Try to give at least one example—or more if you can!

### 3 Extending the Domain of the Logarithm


One of the strongest voices in the effort to expand the scope of the logarithm function was Leonhard Euler (1707–1783), the leading mathematician and physicist of the eighteenth century. He had been thinking about logarithms as early as 1727, in correspondence with his mentor Johann Bernoulli (1667–1748). Bernoulli had written in the past that  $\log(-x) = \log(x)$  (so, in particular,  $\log(-1) = 0$ ). Euler was not convinced, though he recognized that there were arguments in favor of this position. Indeed, in a letter to Bernoulli dated December 10, 1728 he suggested one such argument himself.<sup>1</sup>



If  $\log(xx) = z$ , it will be that

$$\frac{1}{2}z = \log(\sqrt{xx}),$$

but  $\sqrt{xx}$  is both  $-x$  as  $+x$ , therefore  $\frac{1}{2}z$  is  $\log(x)$  and  $\log(-x)$ . But it could be objected that, if  $xx$  has two logarithms, it should be judged to have infinitely many.



**Task 4** Explain Euler’s argument in your own words.

**Task 5** Does Euler’s argument convince you (at this point) that  $\log(-x) = \log(x)$ ? Why or why not?

**Task 6** What argument do you think Euler may have had in mind when he stated that if the logarithmic function has two values, then “it should be judged to have infinitely many”? Write down as many ideas as you can come up with for this. Do you agree with the claim at this point? Why or why not?


### 4 The Beginning of a Conflict

By the end of 1746, Euler was much more mature, both as a mathematician in general and in his understanding of logarithms. Indeed, he had come to strongly suspect that the values of logs of negative values would involve imaginary and complex numbers, although he had yet to publish these ideas. On December 29 of that year, Euler wrote a letter to Jean le Rond d’Alembert (1717–1783), a French philosopher and thinker who had just submitted a paper to the Berlin *Memoires*

---

<sup>1</sup>All translations of excerpts from Euler’s correspondence and other primary sources in this project were prepared by the project author.

(a mathematics journal edited by Euler) in which he tried to prove the Fundamental Theorem of Algebra.<sup>2</sup> In his paper, d’Alembert wrote that for any positive number  $x$ ,  $\log(-x) = \log(x)$ . Euler wrote privately to d’Alembert in part to alert his colleague that this part of his paper may be in error. Regarding the logarithms of negative numbers, Euler wrote:




For me, I believe I have demonstrated [in another work, not yet published] that it [the logarithm of  $-1$ ] is imaginary, and that it is  $= \pi(1 \pm 2n)\sqrt{-1}$ , where  $\pi$  denotes the circumference of a circle in which the diameter  $= 1$ , and  $n$  is any integer.<sup>3</sup>

Because I have shown that, just as each value of sine corresponds to an infinite number of arcs of the circle, so the logarithm of each number has an infinite number of values, among which there is but one that is real when the number is positive, and when the number is negative all the values are imaginary. Thus,  $\log(1) = \pi(0 \pm 2n)\sqrt{-1}$  where  $n$  is any integer. Setting  $n = 0$ , we have the ordinary logarithm  $\log 1 = 0$ . And in the same manner, we have  $\log a = \log a + \pi(0 \pm 2n)\sqrt{-1}$ , where  $\log a$  on the right side of the equation denotes the ordinary logarithm of  $a$ , but  $\log(-a) = \log a + \pi(1 \pm 2n)\sqrt{-1}$ , where all the values are imaginary. All of this follows from the formula

$$\log(\cos \theta + \sin \theta \sqrt{-1})^k = (k\theta \pm 2mk\pi \pm 2n\pi)\sqrt{-1},$$

where  $m$  and  $n$  are any integers, the truth of which is easy to demonstrate.<sup>4</sup>



**Task 7** What did Euler mean when he wrote “corresponds to an infinite number of arcs of the circle”? Use this idea to give an example of a function with infinitely many values for one input. Does this function match the definition given above Task 2?

**Task 8** Assuming that Euler is correct, give three different values of  $\log 1$ .

The next two tasks ask you to investigate Euler’s formula

$$\log(\cos \theta + \sin \theta \sqrt{-1})^k = (k\theta \pm 2mk\pi \pm 2n\pi)\sqrt{-1}.$$

**Task 9** Using this formula, pick values of  $k, m$ , and  $n$  that allow you to calculate  $\log(-1)$ . What do you find?

**Task 10** Now set  $\theta = \pi, n = 0, m = 0$ , and  $k = 1$ . Then exponentiate both sides of the equation and rewrite the equation so that you have 0 on the right hand side, and all other terms on the left. What do you get?

<sup>2</sup>The Fundamental Theorem of Algebra is often stated today as saying that every polynomial of degree  $n$  with complex coefficients has exactly  $n$  complex roots, counting multiplicity. This is equivalent to the claim that every real polynomial has real or complex roots, which is the version that d’Alembert set out to prove.

<sup>3</sup>Euler would eventually use the symbol  $i$  to represent  $\sqrt{-1}$ , but not until 1777.

<sup>4</sup>Euler may have been understating the difficulty in establishing this equation. It didn’t appear until the final page of the paper that he was working on at the time on the logarithms of negative and imaginary numbers. We will take a look at part of this paper in the final section of this project.





## References

- Johann Bernoulli. Solution d'un problème concernant le calcul intégral (Solution of a problem concerning integral calculus). *Histoire de l'Académie Royale des Sciences*, pages 289–297, 1702.
- R. E. Bradley. Euler, D'Alembert and the Logarithm Function. In R. E. Bradley and E. Sandifer, editors, *Leonhard Euler: Life, Work and Legacy*, volume 5 of *Studies in the History and Philosophy of Mathematics*, pages 255–277. Elsevier, Amsterdam, 2007.
- L. Euler. Sur les logarithmes des nombres négatifs et imaginaires (On the logarithms of negative and imaginary numbers). In *Opera Postuma mathematica et physica*, pages 269–281. Imperial Academy of Sciences, St. Petersburg, 1862. Also in *Opera Omnia Series 1*, Volume 19, Birkhäuser, Switzerland, pages 417–438. This posthumously published paper was written by Euler in 1747.
- L. Euler. *Correspondence de Leonhard Euler avec A. C. Clairaut, J. d'Alembert et J. L. Lagrange (Correspondance of Leonhard Euler with A. C. Clairaut, J. d'Alembert and J. L. Lagrange)*. 1980. Volume 5 of Euler's *Opera Omnia*, Series IVA: *Commercium Epistolicum*, eds. A. P. Juškevič and R. Taton, R., Birkhäuser, Basel.
- L. Euler. *Correspondence de Leonhard Euler avec Johann (I) and Niklaus (I) Bernoulli (Correspondance of Leonhard Euler with Johann (I) and Niklaus (I) Bernoulli)*. 1988. Volume 2 of Euler's *Opera Omnia*, Series IVA: *Commercium Epistolicum*, eds. E. A. Fellman and G. K. Mikhajlov, Birkhäuser, Basel.

## Notes to Instructors

### PSP Content: Topics and Goals

This Primary Source Project (PSP) is intended to be used in a course on complex variables, and introduces students to the definition of the logarithm function on negative and complex numbers via an epistolary argument between Leonhard Euler and Jean d’Alembert. Its primary goal is to motivate the definition of the logarithm over  $\mathbb{C}$ , in particular its property of having multiple values. Students familiar with the definition of a function from a real analysis class (or even the vertical line test in calculus) have been taught that a function can have only one output for every input, a lesson belied by the behavior of natural logarithm (along with several other functions) over the complex numbers.

The biggest pedagogical hurdle that this PSP tries to help students clear is thus the idea of a multi-valued function, which might seem a contradiction in terms based on their previous experience. In fact, it was this exact property of the logarithms that made extending the domain of this function to negative and natural numbers so difficult in the eighteenth century. Johann Bernoulli and Jean d’Alembert both seemed unable to make the leap to multi-valued functions, and even Euler (who eventually extended the domain of the logarithm function in the way we think of it today) struggled for several years.

### Student Prerequisites

Ideally, students should have seen a formal definition of a function. They should have a level of mathematical maturity sufficient to look for their own examples in a few places and to be able to reach conclusions using properties of logarithms.

### PSP Design and Task Commentary

Through this project, students have an opportunity to recapitulate some of the early work on the logarithm of negative and imaginary values, and to try to solve an early paradox, proposed by d’Alembert, that seems to indicate that  $\log(-1) = 0$ .

Task 3 can be tricky for students; a multi-valued function doesn’t exist by definition. The hope here is that they may think of  $f(x) = \sqrt{x}$ , which we can think of as a two-valued function, or  $g(x) = \arcsin x$ , which could (through appropriate choices of domain and range) take more. The question of arcsine is explored in Task 7.

Task 7 is designed to help students start thinking about multi-valued functions by reminding them of sine and arcsine. As they know, there are infinitely many angles  $\theta$  for which the  $\sin(\theta) = \pi$ , and thus  $\arcsin(\pi)$  can be thought of as a multi-valued function. We usually avoid this issue and calculus and precalculus classes by restricting the domain and range of arcsine, but one could imagine not doing so.

Task 13 asks students to use a formula for the area of a sector of a circle that leads to an expression involving the logarithm of a complex number. They should find an imaginary value for this logarithm, leading to a real result for the circular sector

## Suggestions for PSP Implementation and Sample Implementation Schedule (based on a 50-minute class period)

One possible implementation, using just 1.5 days of class time, follows. If time allows, the homework can be eliminated and replaced with in-class time, in which case it may take 2–2.5 days to complete the PSP.

- **Day 0.** Students work in groups for about 15 minutes on Day 0 to complete Part 1.

**Day 0 Homework:** Assign Part 2, the reading in Part 3 up to the Task 7, and Task 7. This is the longest section of reading in the PSP, and it may be better done as homework than during class.

- **Day 1.** Students meet in groups to discuss their answers to Task 7, and work together to complete Part 3 and Part 4 through Task 13. The remaining tasks can be completed as homework.

**Day 1 Homework:** Assign Tasks 14–18.

- **Day 2.** I would encourage beginning Day 2 with a short full-class discussion of some of the juicier tasks. Task 17 would be a really good one to work through, and the instructor should feel free to use Task 18 to discuss multiply-valued functions in complex analysis. If it's appropriate for the current state of your course, this could lead to a discussion of branch cuts.

## Connections to other Primary Source Projects

The following primary source-based projects are also freely available for use in teaching courses in complex variables. The number of class periods required for full implementation is given in parentheses. Classroom-ready pdf versions of each can be obtained (along with their  $\text{\LaTeX}$  code) from their authors or downloaded from [https://digitalcommons.ursinus.edu/triumphs\\_complex/](https://digitalcommons.ursinus.edu/triumphs_complex/).

- *Euler's Square Root Laws for Negative Numbers* by David Ruch (1–2 days)
- *An Introduction to Algebra and Geometry in the Complex Plane* by Nicholas A. Scoville and Diana White (5 days)
- *Argand's Development of the Complex Plane* by Nicholas A. Scoville and Diana White (5 days)
- *Gauss and the First "Rigorous" Proof of the Fundamental Theorem of Algebra* by Sarah Hagen and Alan Kappler (3–5 days); also suitable for use in an Introduction to Proofs course
- *Riemann's Development of the Cauchy-Riemann Equations* by David Ruch (3 days)
- *Gauss and Cauchy on Complex Integration* by David Ruch (3 days)


## Recommendations for Further Reading

This project was inspired by Robert Bradley's 2007 article "Euler, D'Alembert and the Logarithm Function." While I have tried to encapsulate the most important ideas from that article in this Primary Source Project, a lot of rich discussion about eighteenth-century views of logarithms were left out. Instructors (and interested students!) are strongly encouraged to read Bradley's paper.



## Acknowledgments

The development of this student project has been partially supported by the TRansforming Instruction in Undergraduate Mathematics via Primary Historical Sources (TRIUMPHS) Program with funding from the National Science Foundation’s Improving Undergraduate STEM Education Program under Grant Nos. 1523494, 1523561, 1523747, 1523753, 1523898, 1524065, and 1524098. Any opinions, findings, and conclusions or recommendations expressed in this project are those of the author and do not necessarily represent the views of the National Science Foundation.



With the exception of excerpts taken from published translations of the primary sources used in this project and any direct quotes from published secondary sources, this work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License (<https://creativecommons.org/licenses/by-sa/4.0/legalcode>). It allows re-distribution and re-use of a licensed work on the conditions that the creator is appropriately credited and that any derivative work is made available under “the same, similar or a compatible license.”

For more information about TRIUMPHS, visit <https://blogs.ursinus.edu/triumphs/>.