



A Calculation of $\int_0^1 e^{-x^2} dx$

Alberto L. Delgado

The College Mathematics Journal, Vol. 34, No. 4. (Sep., 2003), pp. 321-323.

Stable URL:

<http://links.jstor.org/sici?sici=0746-8342%28200309%2934%3A4%3C321%3AACO%3E2.0.CO%3B2-M>

The College Mathematics Journal is currently published by Mathematical Association of America.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.


Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact support@jstor.org.

References

1. W. A. McGeveran Jr., editor, *The World Almanac and Book of Facts 2002*, World Almanac Education Group, New York, 2002.
2. A. E. Roy, *Orbital Motion*, 3rd ed., Adam Hilger, Ltd., Bristol, UK, 1988.
3. R. Urschel, *Analemma*, <http://www.analemma.com>.



A Calculation of $\int_0^\infty e^{-x^2} dx$

Alberto L. Delgado (delgado@hilltop.bradley.edu) Bradley University, Peoria, IL 61625-0160

If you are looking for a way to tie together different strands of material your students see in their second semester of a standard three-semester calculus course, the result in the title will serve you very well. Of course, the result is

$$\int_0^\infty e^{-x^2} dx = \frac{\sqrt{\pi}}{2}.$$

Students find it surprising (where does the π come from?) and meaningful (when they learn its relationship to the bell curve). You can also use this significant improper integral as an entry into a discussion of approximations of definite integrals over finite regions.

There are numerous ways to evaluate the integral in the title. In just a few minutes in the library, I located two evaluations based on Wallis' infinite product expansion of π , [8] and [5]; a calculation using contour integration in the complex plane, [1]; a reduction to Euler's integral of the first kind, [2]; two evaluations using differentiation under the integral sign, [7] and [9]; and a calculation based on solids of revolution, [3]. You can find the standard computation based on a double integral in polar coordinates in almost any statistics or calculus book. The integral is commonly associated with Gauss, although he credits Laplace with its discovery and publication in 1805. Euler was working with similar integrals thirty years earlier, but he seems to have missed its exact formulation. By 1813, the result appears to have been well known. See [6] for a more detailed history.

Few of the techniques for evaluating this integral are easily accessible to a beginning student. Here is a technique that students can appreciate. Although the method may not be new (the main idea is already in [3] and [4]) our variation of it is sufficiently elementary that it can be presented to students.

Let I denote the value of the integral of interest. A comparison with e^{-x} confirms that I is finite. Let's start by finding the volume V of the solid of revolution made by revolving the graph of e^{-x^2} for $x \in [0, \infty)$ about the y -axis (Figure 1).

Using the usual technique of nested cylindrical shells, we obtain

$$V = \int_0^\infty 2\pi x e^{-x^2} dx = \pi.$$

Now let's compute V a second time. This time take cross sections parallel to the x -axis (Figure 2).




Figure 1. Graph of e^{-x^2} rotated about y -axis.




Figure 2. Cross section at distance t from xy -plane.

Denoting the area of the cross section at distance t from the x -axis by A_t , we get $V = 2 \int_0^\infty A_t dt$. In order to compute A_t , first find the equation for the graph of the cross section. Consider the point P on the cross section. Since the surface is radially symmetric, we can find P 's height by rotating back to the value $d = \sqrt{u^2 + t^2}$ on the x -axis. Thus, P 's height is $e^{-(\sqrt{u^2+t^2})^2}$, and

$$A_t = 2 \int_0^\infty e^{-(u^2+t^2)} du = 2 \int_0^\infty e^{-u^2} e^{-t^2} du = 2e^{-t^2} \int_0^\infty e^{-u^2} du = 2e^{-t^2} I.$$

Therefore,


$$V = 2 \int_0^\infty A_t dt = 2 \int_0^\infty 2e^{-t^2} I dt = 4I \int_0^\infty e^{-t^2} dt = 4I^2.$$

Equating the two expressions for V , we obtain $\pi = 4I^2$ and the result follows.

References

1. Darrell Desbrow, On evaluating $\int_{-\infty}^\infty e^{ax(x-2b)} dx$ by contour integration round a parallelogram, *American Mathematical Monthly* **105** (1998) 726–731.

2. Édouard Goursat, *A course in mathematical analysis, vol. I*, Ginn & Co., 1904.
3. C. P. Nicholas, Another look at the probability integral, *American Mathematical Monthly* **64** (1957) 739–741.
4. C. P. Nicholas and R. C. Yates, The probability integral, *American Mathematical Monthly* **57** (1950) 412–413.
5. T. J. Stieljes, *Oevres Complètes, vol. II*, Wolters-Noordhoff, 1918.
6. S. C. van Veen, Historische bijzonderheden over $\int_0^\infty e^{-x^2} dx$, *Mathematica B* **12** (1943–1944), 1–4.
7. Robert Weinstock, Elementary evaluations of $\int_0^\infty e^{-x^2} dx$, $\int_0^\infty \cos(x^2) dx$, and $\int_0^\infty \sin(x^2) dx$, *American Mathematical Monthly* **97** (1990) 39–42.
8. Robert M. Young, *Excursions in Calculus*, Mathematical Association of America, 1992.
9. ———, On the evaluation of certain improper integrals, *Mathematical Gazette* **75** (1991) 88–89.



A Simple Introduction to e

Pratibha Ghatage (ghatage@math.csuohio.edu) Cleveland State University, Cleveland, OH 44115

The idea for this note was suggested by Dimitric’s recent paper [1] that discusses a way to introduce the Euler number e in a heuristic way. Given the volume and breadth of literature on the subject, what follows is a very simplistic way of introducing e that is easily understood by freshmen.

Let A_2 denote the area under the hyperbola $y = \frac{1}{x}$ between $x = 1$ and $x = 2$. Then (Figure 1) A_2 is greater than the area $\frac{1}{2}$ of the “inner” rectangle having height $\frac{1}{2}$, and




Figure 1.




Figure 2.




Figure 3.

LINKED CITATIONS

- Page 1 of 2 -



You have printed the following article:

A Calculation of $\int_0^x e^{-x^2} dx$

Alberto L. Delgado

The College Mathematics Journal, Vol. 34, No. 4. (Sep., 2003), pp. 321-323.

Stable URL:

<http://links.jstor.org/sici?sici=0746-8342%28200309%2934%3A4%3C321%3AACO%3E2.0.CO%3B2-M>

This article references the following linked citations. If you are trying to access articles from an off-campus location, you may be required to first logon via your library web site to access JSTOR. Please visit your library's website or contact a librarian to learn about options for remote access to JSTOR.

References

¹ **On Evaluating $\int_{-a}^a e^{ax(x-2b)} dx$ by Contour Integration Round a Parallelogram**

Darrell Desbrow

The American Mathematical Monthly, Vol. 105, No. 8. (Oct., 1998), pp. 726-731.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28199810%29105%3A8%3C726%3A0EBCIR%3E2.0.CO%3B2-P>

³ **Another Look at the Probability Integral**

C. P. Nicholas

The American Mathematical Monthly, Vol. 64, No. 10. (Dec., 1957), pp. 739-741.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28195712%2964%3A10%3C739%3AALATPI%3E2.0.CO%3B2-L>

⁴ **The Probability Integral**

C. P. Nicholas; R. C. Yates

The American Mathematical Monthly, Vol. 57, No. 6. (Jun. - Jul., 1950), pp. 412-413.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28195006%2F07%2957%3A6%3C412%3ATPI%3E2.0.CO%3B2-4>

⁷ **Elementary Evaluations of $\int_0^{\infty} e^{-x^2} dx$, $\int_0^{\infty} \cos x^2 dx$ and $\int_0^{\infty} \sin x^2 dx$**

Robert Weinstock

The American Mathematical Monthly, Vol. 97, No. 1. (Jan., 1990), pp. 39-42.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28199001%2997%3A1%3C39%3AEEOA%3E2.0.CO%3B2-4>

NOTE: *The reference numbering from the original has been maintained in this citation list.*

LINKED CITATIONS

- Page 2 of 2 -



⁹ **75.4 On the Evaluation of Certain Improper Integrals**

Robert M. Young

The Mathematical Gazette, Vol. 75, No. 471. (Mar., 1991), pp. 88-89.

Stable URL:

<http://links.jstor.org/sici?sici=0025-5572%28199103%292%3A75%3A471%3C88%3A7OTE0C%3E2.0.CO%3B2-P>