



The Spider's Spacewalk Derivation of \sin' and \cos'

Tim Hesterberg

The College Mathematics Journal, Vol. 26, No. 2. (Mar., 1995), pp. 144-145.

Stable URL:

<http://links.jstor.org/sici?sici=0746-8342%28199503%2926%3A2%3C144%3ATSSDOS%3E2.0.CO%3B2-9>

The College Mathematics Journal is currently published by Mathematical Association of America.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact support@jstor.org.

The Spider's Spacewalk Derivation of \sin' and \cos

Tim Hesterberg, Franklin and Marshall College, Lancaster, PA 17604-3003

The usual proofs of the derivatives of sine and cosine in introductory calculus involve limits. I shall outline a simple geometric derivation that avoids evaluating limits, based on the interpretation of the derivative as the instantaneous rate of change. The principle behind this proof is found in a late nineteenth-century calculus textbook by J. M. Rice and W. W. Johnson, *The Elements of the Differential Calculus, Founded on the Method of Rates or Fluxions* (Wiley, New York, 1874).

A spider walks with speed 1 in a circular path around the outside of a round satellite of radius 1, as shown in Figure 1. At time t the spider will have travelled a distance t , which corresponds to a central angle of t radians. The altitude of the spider, in the standard coordinate system, is $y = \sin(t)$ and the spider is $x = \cos(t)$ units to the right (or left) of the origin.

Now look how fast the spider is moving *upward*. Since the altitude of the spider at time t is $y = \sin(t)$, its upward velocity is $y' = \sin'(t)$. Oops!—The spider loses its footing at time t , and since the gravity in outer space is negligible, it continues with the same direction and speed. It moves a distance 1 in additional time $\Delta t = 1$,




Figure 1




Figure 2

from point S to point B ; see Figure 2. The spider's altitude changes by $\Delta y = SC$, so its upward velocity is $y' = \Delta y / \Delta t = SC$. But because triangles OAS and SCB are congruent ($\angle OSA = 90^\circ - t$, so $\angle CSB = t = \angle AOS$), $SC = OA = x = \cos(t)$. Therefore $\sin'(t) = y' = \cos(t)$.

Similarly, the spider is $x = \cos(t)$ units to the right of the center, and its horizontal velocity is $x' = \cos'(t) = \Delta x / \Delta t = -BC = -SA = -\sin(t)$. The minus sign arises because the spider is moving to the left when it is above the x -axis, i.e., $\Delta x / \Delta t$ is negative whenever $y = \sin(t)$ is positive.


Figure 2 describes the case $0 < t < \pi/2$, but it is easily modified to yield the same results when the spider is located anywhere on the unit circle.

Acknowledgment. I thank Bob Gether and George Rosenstein for helpful comments and for the reference to Rice and Johnson's book.

[*Editor's note.* This proof also appeared in C. S. Ogilvy, Derivatives of $\sin \theta$ and $\cos \theta$, *American Mathematical Monthly* 67 (1960) 673.]

—o—

$\cos(x+y)$ for $x+y > \pi/2$



$$\begin{aligned} \frac{1}{2}ab\sin[\pi/2 - (x+y)] &= \frac{1}{2}b \cos x \cdot a \cos y - \frac{1}{2}a \sin y \cdot b \sin x \\ \therefore \cos(x+y) &= \cos x \cdot \cos y - \sin x \cdot \sin y \end{aligned}$$

Sidney H. Kung